



SHILAP Revista de Lepidopterología

ISSN: 0300-5267

avives@orange.es

Sociedad Hispano-Luso-Americana de  
Lepidopterología  
España

Efetov, K. A.; Kucherenko, E. E.; Parshkova, E. V.; Tarmann, G. M.  
2-butyl 2-dodecenoate, a new sex attractant for *Jordanita* (*Tremewania*) *notata* (Zeller,  
1847) and some other Procrinae species (Lepidoptera: Zygaenidae)  
SHILAP Revista de Lepidopterología, vol. 44, núm. 175, septiembre, 2016, pp. 519-527  
Sociedad Hispano-Luso-Americana de Lepidopterología  
Madrid, España

Available in: <http://www.redalyc.org/articulo.oa?id=45549999013>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## **2-butyl 2-dodecenoate, a new sex attractant for *Jordanita (Tremewania) notata* (Zeller, 1847) and some other Procridinae species (Lepidoptera: Zygaenidae)**

K. A. Efetov, E. E. Kucherenko, E. V. Parshkova &  
G. M. Tarmann

### **Abstract**

A property of 2-butyl 2-dodecenoate as a sex attractant for the males of *Jordanita (Tremewania) notata* (Zeller, 1847) was proven in field trapping tests and behavioural observations in the Crimean Peninsula. It was shown that this substance is also attractive for males of some other Procridinae species, viz. *Rhagades (Rhagades) pruni* ([Denis & Schiffermüller], 1775), *Adscita (Adscita) geryon* (Hübner, 1813), *Jordanita (Jordanita) graeca* (Jordan, 1907), *J. (J.) globulariae* (Hübner, 1793), and *J. (Solani) subsolana* (Staudinger, 1862). The attractiveness of 2-butyl 2-dodecenoate for *Rh. pruni*, *A. geryon*, *J. notata*, and *J. subsolana* was found for the first time.

KEY WORDS: Lepidoptera, Zygaenidae, Procridinae, *Jordanita notata*, *J. graeca*, *J. globulariae*, *J. subsolana*, *Adscita geryon*, *Rhagades pruni*, 2-butyl 2-dodecenoate, sex attractant, Crimea.

**2-butyl 2-dodecenoato, un nuevo atrayente sexual para *Jordanita (Tremewania) notata*  
(Zeller, 1847) y algunas otras especies de Procridinae  
(Lepidoptera: Zygaenidae)**

### **Resumen**

2-butyl 2-dodecenoato ha sido sintetizado y sus propiedades como atrayente sexual para los machos de *Jordanita (Tremewania) notata* (Zeller, 1847) fueron probadas en trampas de campo y observado el comportamiento en la Península de Crimea. Se ha demostrado que esta sustancia también es atractiva para los machos de algunas otras especies de Procridinae, véase *Rhagades (Rhagades) pruni* ([Denis & Schiffermüller], 1775), *Adscita (Adscita) geryon* (Hübner, 1813), *Jordanita (Jordanita) graeca* (Jordan, 1907), *J. (J.) globulariae* (Hübner, 1793) y *J. (Solani) subsolana* (Staudinger, 1862).

PALABRAS CLAVE: Lepidoptera, Zygaenidae, Procridinae, *Jordanita notata*, *J. graeca*, *J. globulariae*, *J. subsolana*, *Adscita geryon*, *Rhagades pruni*, 2-butyl 2-dodecenoato, atrayente sexual, Crimea.

### **Introduction**

Currently, sex pheromones of insects are widely used for the detection and direct control of species (both protected and pest species) and for monitoring their numbers in various biotopes (MILLAR *et al.*, 2010; OLEANDER *et al.*, 2015; SUBCHEV, 2014; WITZGALL *et al.*, 2010). Most probably, female sex pheromones are the main factor in the attraction and recognition of females by conspecific males in some groups of Lepidoptera that are represented by habitually very

similar species, for example in the subfamily Procrinae of the family Zygaenidae (EFETOV, 1996, 1997a, 1997b, 1998, 2001a, 2001c, 2006, 2010; EFETOV *et al.*, 2004; EFETOV & TARMANN, 2013a, 2013b, 2014a, 2014b; TOSHOVA *et al.*, 2007; ZAGATTI & RENOU, 1984). The Procrinae are divided into two tribes: Artonini Tarmann, 1994, and Procradini Boisduval, 1828 (TARMANN, 1994; EFETOV *et al.*, 2000, 2006; EFETOV & HAYASHI, 2008). The Procrinae are represented only by the Procradini in the western Palaearctis (EFETOV, 2005; EFETOV & TARMANN, 2012).

One Procradini species, *Jordanita (Tremewania) notata* (Zeller, 1847), is distributed from Spain and Portugal through central and southern Europe to Ukraine, Crimea, Northern Caucasus, Transcaucasia, Turkey, and north-western Iran (EFETOV & TARMANN, 1999; EFETOV, 2004, 2005). Not long ago it was shown that (2R)-butyl (7Z)-dodecenoate, previously established as a female sex pheromone component of *Illiberis (Primilliberis) rotundata* Jordan, 1907 (SUBCHEV *et al.*, 2009), is a sex attractant for the males of *J. notata* in the Crimea, Bulgaria (SUBCHEV *et al.*, 2010) and Italy (EFETOV *et al.*, 2012, 2015). In Eurasia the enantiomers of 2-butyl (7Z)-dodecenoate and/or their mixtures are also sex attractants for males of some other Procradini species, viz. *Rhagades (Rhagades) pruni* ([Denis & Schiffermüller], 1775), *Adscita (Adscita) geryon* (Hübner, 1813), *Adscita (Tarmannita) mannii* (Lederer, 1853) (SUBCHEV *et al.*, 2010; EFETOV *et al.*, 2012), *Zygaenoprocris (Molletia) taftana* (Alberti, 1939) (EFETOV *et al.*, 2011), *Z. (Zygaenoprocris) eberti* (Alberti, 1968) (EFETOV *et al.*, 2014a), *J. (Praviela) anatolica* (Naufock, 1929) (EFETOV *et al.*, 2010), and *J. (Rjabovia) horni* (Alberti, 1937) (EFETOV *et al.*, 2011). The mixtures of (2R)-butyl (7Z)-dodecenoate and (2R)-butyl (9Z)-tetradecenoate are sex attractants for males of *I. (P.) rotundata* (SUBCHEV *et al.*, 2012) and *I. (P.) pruni* Dyar, 1905 (SUBCHEV *et al.*, 2013).

The aim of this work was to study the biological activity of another ester of sec-butanol and dodecenoate with a double bond in the second position in dodecenoate. This substance has been synthesized in our laboratory and its attractiveness for different species of the Procrinae was tested during field observations in the Crimea in 2013-2014.

## Materials and methods

Sec-butanol (Sigma-Aldrich, Germany) and other reagents (Ukraine) were used for the synthesis of 2-butyl 2-dodecenoate (EFETOV *et al.*, 2013, 2014b). The products of reactions were separated by vacuum distillation, the content of products was determined by a gas-liquid chromatography with a chromatograph "Tsvet-500", thermal conductivity detector (katharometer). The column of 3 m, SP-2250 on Supelcoports, temperature of the column is 90-130° C, of the evaporator - 130-220° C, of the detector - 90-190° C. A chemical nature of the synthesized substance was confirmed using a method of nuclear magnetic resonance spectroscopy (EFETOV *et al.*, 2014b).

For preparing baits, 2-butyl 2-dodecenoate was applied onto vial caps (200 microlitres per cap without solvent) composed of grey rubber. The baits were tested in home-made sticky Delta traps with removable sticky layers covered with Tanglefoot® insect glue and hung on bushes or trees at a height of 1.0-1.5 m above the ground. In all sites we also placed control traps with rubber caps but without the attractant (distance to the baited traps was not less than 10 metres).

Sometimes we merely placed the lure on stones on the ground. In this case we collected attracted specimens by netting them near the lure.

Field observations were made in six localities in the mountain region of the Crimean Peninsula (Fig. 1). A list of studied localities and periods of observation is provided below.

SITE I: vic. Dachnoye, N of Sudak, 110 m. Bushy mountain slope near deciduous forest. Periods of observation: 16-VI-2013 - 18-VII-2013, one baited trap and one control trap; 3-V-2014 - 27-VII-2014, one baited trap and one control trap.

SITE II: vic. Belogorsk, Mt. Sary-Kaya, 230-239 m. Grassy slope near steep calcareous edge of

the mountain. Periods of observation: 30-V-2013 – 27-VII-2013, two baited traps and one control trap; 3-V-2014 - 27-VII-2014, one baited trap and one control trap.

SITE III: vic. Simferopol, Bitak, 320 m. Grassy slope near steep calcareous edge of the mountain. Periods of observation: 28-IV-2013 - 27-VII-2013, two baited traps and one control trap; 13-V-2014 - 31-VII-2014, one baited trap and one control trap.


SITE IV: Mt. Chatyr-Dag, 466-514 m. Clearings in deciduous forest. Periods of observation: 5-VI-2013 - 4-VIII-2013, one baited trap and one control trap; 17-V-2014 - 31-VII-2014, three baited traps and two control traps.

SITE V: vic. Izobilnoye, N of Alushta, 180 m. Bushy slope between a vineyard and deciduous forest. Periods of observation: 4-V-2014 - 7-VIII-2014, two baited traps and one control trap.

SITE VI: vic. Luchistoye, NE of Alushta, 376-388 m. Bushy mountain slope near the lake. Periods of observation: 4-V-2014 - 7-VIII-2014, two baited traps and one control trap.

The nomenclature of the subfamily Procridinae follows the last revisions of the group (EFETOV, 2001b, 2001c; EFETOV & TARMANN, 2012).

All collected specimens have been determined by K. A. Efetov, based on examination of the genitalia.



**Figure 1.**– Map of the Crimea showing the sites that were investigated using traps baited with 2-butyl 2-dodecenoate. 1. vic. Dachnoye, N of Sudak; 2. vic. Belogorsk, Mt. Sary-Kaya; 3. vic. Simferopol, Bitak; 4. Mt. Chatyr-Dag; 5. vic. Izobilnoye, N of Alushta; 6. vic. Luchistoye, NE of Alushta.

## Results and discussion

The total number of specimens attracted is shown in Table 1. Dates of inspection of the traps in different biotopes and type of attraction of specimens (glued in trap or netted near lures) are listed below. Procridinae males were absent in control traps (without attractant) in all localities.

**Table 1.**— Total number of males of six Procridinae species attracted by traps baited with 2-butyl 2-dodecenoate in the Crimea. In all localities specimens were absent in control traps (without attractant).

Locality	vic. Dachnoye (site I)	vic. Belogorsk (site II)		vic. Simferopol (site III)		Mt. Chatyr-Dag (site IV)		vic. Izobilnoye (site V)	vic. Luchistoye (site VI)	Total number of males
Species	2014	2013	2014	2013	2014	2013	2014	2014	2014	
<i>Rh. pruni</i>									2	2
<i>A. geryon</i>					8					8
<i>J. notata</i>		*	125							125
<i>J. graeca</i>	1	36	10	1				1	46	95
<i>J. globulariae</i>						5	5			10
<i>J. subsolana</i>								1	2	3

\* - In 2013 the traps were placed in site II after the flight period of *J. notata* had already finished.

*Rhagades (Rhagades) pruni* ([Denis & Schiffermüller], 1775)

vic. Luchistoye, NE of Alushta, 376 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 14-VI-2014; 1 ♂, 21-VI-2014.

*Adscita (Adscita) geryon* (Hübner, 1813)

vic. Simferopol, Bitak, 320 m, near lure with 2-butyl 2-dodecenoate: 4 ♂♂, 17-V-2014; 4 ♂♂, 27-V-2014.

*Jordanita (Tremewanina) notata* (Zeller, 1847)

vic. Belogorsk, Mt. Sary-Kaya, 239 m, near lure with 2-butyl 2-dodecenoate: 1 ♂, 3-V-2014; in trap with 2-butyl 2-dodecenoate: 51 ♂♂, 18-V-2014; 17 ♂♂, 24-V-2014; 56 ♂♂, 1-VI-2014.

*Jordanita (Jordanita) graeca* (Jordan, 1907)

vic. Dachnoye, N of Sudak, 110 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 1-VI-2014. Vic. Belogorsk, Mt. Sary-Kaya, 230 m, near lure with 2-butyl 2-dodecenoate: 7 ♂♂, 30-V-2013; 26 ♂♂, 11-VI-2013, 2 ♂♂, 19-VI-2013; 10 ♂♂, 24-V-2014; in trap with 2-butyl 2-dodecenoate: 1 ♂, 11-VI-2013. Vic. Simferopol, Bitak, 320 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 24-V-2013. Vic. Izobilnoye, N of Alushta, 180 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 25-V-2014. Vic. Luchistoye, NE of Alushta, 376 m, near lure with 2-butyl 2-dodecenoate: 2 ♂♂, 29-V-2014; in trap with 2-butyl 2-dodecenoate: 1 ♂, 25-V-2014; 388 m, near lure with 2-butyl 2-dodecenoate: 21 ♂♂, 25-V-2014; 21 ♂♂, 29-V-2014; 1 ♂, 31-V-2014.

*Jordanita (Jordanita) globulariae* (Hübner, 1793)

Mt. Chatyr-Dag, 466 m, in trap with 2-butyl 2-dodecenoate: 5 ♂♂, 23-VI-2013; 1 ♂, 31-V-2014; 2 ♂♂, 14-VI-2014; 514 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 14-VI-2014; 1 ♂, 28-VI-2014.

*Jordanita (Solaniterna) subsolana* (Staudinger, 1862)

vic. Izobilnoye, N of Alushta, 180 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 29-V-2014. Vic. Luchistoye, NE of Alushta, 376 m, in trap with 2-butyl 2-dodecenoate: 1 ♂, 25-V-2014; 1 ♂, 29-V-2014.

An application of 2-butyl 2-dodecenoate demonstrated its high attractiveness for the males of *J. notata*. 125 male specimens were collected near Belogorsk from 3-V-2014 to 27-VII-2014, 124 of which were found in one sticky trap that was controlled once per week. The number of attracted males could have been higher, as the sticky layers during the maximum flight period every week were completely covered with specimens (Fig. 2).



**Figure 2.**– Sticky trap baited with 2-butyl 2-dodecenoate with 56 males of *J. notata*, vic. Belogorsk, Mt. Sary-Kaya, 1-VI-2014.



Except for *J. notata*, males of *Rh. pruni*, *A. geryon*, *J. graeca*, *J. globulariae*, and *J. subsolana* were also attracted during our field observations. We found two different types of attraction.

1. *J. notata* and *J. globulariae* were found as glued specimens on sticky layers. *Rh. pruni* and *J. subsolana* were also found in sticky traps but in small numbers (two and three specimens respectively).

2. The other two species, viz. *J. graeca* and *A. geryon*, came actively to the lures (rubber caps with 2-butyl 2-dodecenoate) that were placed in the biotope. Only five specimens (out of 95) of *J. graeca* were found in sticky traps. These differences in attraction are shown in Figs 3 and 4. Most probably the males (that came to the lure and did not adhere to the sticky layers) realized at a close distance that the 2-butyl 2-dodecenoate is not their natural pheromone.

## Conclusion

It was shown that synthetic 2-butyl 2-dodecenoate attracts the males of *Jordanita (Tremewania) notata* and other Procridinae species: *Rhagades (Rhagades) pruni*, *Adscita (Adscita) geryon*, *Jordanita (Jordanita) graeca*, *J. (J.) globulariae*, and *J. (Solaniterna) subsolana*. This substance can be used for identifying the presence of Procridinae species in different regions and for seasonal monitoring of these moths. While the attraction of *J. graeca* and *J. globulariae* by 2-butyl 2-dodecenoate we have already registered earlier (EFETOV *et al.*, 2014b), the attractiveness of this substance for *Rh. pruni*, *A. geryon*, *J. notata*, and *J. subsolana* was found for the first time.



**Figures 3-4.**– 3. Attraction of *Jordanita notata* by 2-butyl 2-dodecenoate. 4. Attraction of *Jordanita graeca* by 2-butyl 2-dodecenoate.



## Acknowledgements

For help in preparing 2-butyl 2-dodecenoate we thank Dr M. Y. Baevsky and Mr A. I. Poddubov (both - Simferopol, Crimea).

## BIBLIOGRAPHY

- EFETOV, K. A., 1996.– The description of the female of *Illiberis (Alterasvenia) yuennanensis* Alberti, 1951 (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **47**(2): 111-113.
- EFETOV, K. A., 1997a.– Two new species of the genus *Artona* Walker, 1854 (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **48**(3): 165-177.
- EFETOV, K. A., 1997b.– Three new species of the genus *Illiberis* Walker, 1854, from Taiwan and Vietnam (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **48**(4): 231-244.
- EFETOV, K. A., 1998.– A revision of the genus *Goe* Hampson, [1893] (Lepidoptera: Zygaenidae, Procridinae), with descriptions of two new species.– *Entomologist's Gazette*, **49**(1): 49-62.
- EFETOV, K. A., 2001a.– On the systematic position of *Zygaenoprocris* Hampson, 1900 (Lepidoptera: Zygaenidae, Procridinae) and the erection of two new subgenera.– *Entomologist's Gazette*, **52**(1): 41-48.
- EFETOV, K. A., 2001b.– An annotated check-list of Forester moths (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **52**(3): 153-162.
- EFETOV, K. A., 2001c.– *A review of the western Palaearctic Procridinae (Lepidoptera: Zygaenidae)*: 328 pp. CSMU Press, Simferopol.
- EFETOV, K. A., 2004.– *Forester and Burnet moths (Lepidoptera: Zygaenidae). The genera Theresimima Strand, 1917, Rhagades Wallengren, 1863, Zygaenoprocris Hampson, 1900, Adscita Retzius, 1783, Jordanita Verity, 1946 (Procridinae), and Zygaena Fabricius, 1775 (Zygaeninae)*: 272 pp. CSMU Press, Simferopol.
- EFETOV, K. A., 2005.– *The Zygaenidae (Lepidoptera) of the Crimea and other regions of Eurasia*: 420 pp. CSMU Press, Simferopol.
- EFETOV, K. A., 2006.– Nine new species of the genus *Chrysartona* Swinhoe, 1892 (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **57**(1): 23-50.
- EFETOV, K. A., 2010.– *Illiberis (Hedina) louisi* sp. nov. (Lepidoptera: Zygaenidae, Procridinae) from China.– *Entomologist's Gazette*, **61**(4): 235-241.
- EFETOV, K. A., BAEVSKY, M. Y., BEKETOV, A. A., PARSHKOVA, E. V. & PODDUBOV, A. I., 2013.– Synthesis of 2-butyl 2-dodecenoate, a possible sex attractant for the species of the subfamily Procridinae (Lepidoptera: Zygaenidae).– *Tavricheskiy Mediko-biologicheskii Vestnik*, **16**(4): 53-57.
- EFETOV, K. A., CAN, F., TOSHOVA, T. B. & SUBCHEV, M., 2010.– New sex attractant for *Jordanita anatolica* (Naufock) (Lepidoptera: Zygaenidae: Procridinae).– *Acta Zoologica Bulgarica*, **62**(2): 315-319.
- EFETOV, K. A. & HAYASHI, E., 2008.– On the chaetotaxy of the first instar larva of *Artona martini* Efetov, 1997 (Lepidoptera: Zygaenidae, Procridinae, Artonini).– *Entomologist's Gazette*, **59**(2): 101-104.
- EFETOV, K. A., HOFMANN, A. & TARMANN, G. M., 2014a.– Application of two molecular approaches (use of sex attractants and DNA barcoding) allowed to rediscover *Zygaenoprocris eberti* (Alberti, 1968) (Lepidoptera, Zygaenidae, Procridinae), hitherto known only from the female holotype.– *Nota Lepidopterologica*, **37**(2): 151-160.
- EFETOV, K. A., KEIL, T., MOLLET, B. & TARMANN, G. M., 2000.– New data on the chaetotaxy of the first instar larva of Forester moths (Lepidoptera: Zygaenidae, Procridinae).– *Nachrichten des Entomologischen Vereins Apollo, Frankfurt am Main, N. F.*, **21**(2): 83-90.
- EFETOV, K. A., PARSHKOVA, E. V., BAEVSKY, M. Y. & PODDUBOV, A. I., 2014b.– Sec-butyl ester of dodecenoate: synthesis and attractive properties.– *The Ukrainian Biochemical Journal*, **86**(6): 175-182.
- EFETOV, K. A., PARSHKOVA, E. V. & KOSHIO C., 2004.– The karyotype of *Illiberis (Primilliberis) rotundata* Jordan, [1907] (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **55**(3): 167-170.
- EFETOV, K. A., SUBCHEV, M. A., TOSHOVA, T. B. & KISELEV, V. M., 2011.– Attraction of *Zygaenoprocris taftana* (Alberti, 1939) and *Jordanita horni* (Alberti, 1937) (Lepidoptera: Zygaenidae, Procridinae) by synthetic sex pheromones in Armenia.– *Entomologist's Gazette*, **62**(2): 113-121.
- EFETOV, K. A. & TARMANN, G. M., 1999.– *Forester moths. The genera Theresimima Strand, 1917,*



- Rhagades Wallengren, 1863, Jordanita Verity, 1946, and Adscita Retzius, 1783 (Lepidoptera: Zygaenidae, Procridinae)*: 192 pp. Apollo Books, Stenstrup.
- EFETOV, K. A. & TARMANN, G. M., 2012.– A checklist of the Palearctic Procridinae (Lepidoptera: Zygaenidae): 108 pp. CSMU Press, Simferopol - Innsbruck.
- EFETOV, K. A. & TARMANN, G. M., 2013a.– *Illiberis (Alterasvenia) cernyi* sp. nov. (Lepidoptera: Zygaenidae, Procridinae) from northern Thailand.– *Entomologist's Gazette*, **64**(1): 33-39.
- EFETOV, K. A. & TARMANN, G. M., 2013b.– *Chrysartona (Chrystarmanna) mineti* sp. nov. (Lepidoptera: Zygaenidae, Procridinae) from northern Vietnam.– *Entomologist's Gazette*, **64**(3): 197-206.
- EFETOV, K. A. & TARMANN, G. M., 2014a.– *Illiberis (Alterasvenia) banmauka* sp. nov. (Lepidoptera: Zygaenidae, Procridinae) from China and Myanmar.– *Entomologist's Gazette*, **65**(1): 62-70.
- EFETOV, K. A. & TARMANN, G. M., 2014b.– A new European species, *Adscita dujardini* sp. nov. (Lepidoptera: Zygaenidae, Procridinae) confirmed by DNA analysis.– *Entomologist's Gazette*, **65**(3): 179-200.
- EFETOV, K. A., TARMANN, G. M., HAYASHI, E. & PARSHKOVA, E. V., 2006.– New data on the chaetotaxy of the first instar larvae of Procridini and Artonini (Lepidoptera: Zygaenidae, Procridinae).– *Entomologist's Gazette*, **57**(4): 229-233.
- EFETOV, K. A., TARMANN, G. M., TOSHOVA, T. B. & SUBCHEV, M. A., 2012.– Attraction of *Adscita mannii* (Lederer, 1853), *A. geryon* (Hübner, 1813) and *Jordanita notata* (Zeller, 1847) (Lepidoptera: Zygaenidae, Procridinae) to 2-butyl 7Z-dodecenoate in Italy.– *XIII International Symposium on Zygaenidae (Innsbruck, 16-23 September 2012)*, **2012**: 15.
- EFETOV, K. A., TARMANN, G. M., TOSHOVA, T. B. & SUBCHEV, M. A., 2015.– Enantiomers of 2-butyl 7Z-dodecenoate are sex attractants for males of *Adscita mannii* (Lederer, 1853), *A. geryon* (Hübner, 1813), and *Jordanita notata* (Zeller, 1847) (Lepidoptera: Zygaenidae, Procridinae) in Italy.– *Nota Lepidopterologica*, **38**(2): 161-169.
- MILLAR, J. G., MCELFRESH, J. S., ROMERO, C., VILA, M., MARI-MENA, N. & LOPEZ-VAAMONDE C., 2010.– Identification of the sex pheromone of a protected species, the Spanish Moon Moth *Graellsia isabellae* (Lepidoptera: Saturniidae).– *Journal of Chemical Ecology*, **36**: 923-932.
- OLEANDER, A., THACKERY, D. & BURMAN, J., 2015 (in press). The effect of exposure to synthetic pheromone lures on male *Zygaena filipendulae* mating behaviour: implications for monitoring species of conservation interest.– *Journal of Insect Conservation*, DOI 10.1007/s10841-015-9775-4
- SUBCHEV, M., 2014.– Sex pheromone communication in the family Zygaenidae (Insecta: Lepidoptera): a review.– *Acta Zoologica Bulgarica*, **66**(2): 147-157.
- SUBCHEV, M., EFETOV, K. A., TOSHOVA, T., PARSHKOVA, E. V., TÓTH, M. & FRANCKE, W., 2010.– New sex attractants for species of the zygaenid subfamily Procridinae (Lepidoptera: Zygaenidae).– *Entomologia Generalis* (Stuttgart), **32**(4): 243-250.
- SUBCHEV, M. A., KOSHIO, C., TOSHOVA, T. B. & EFETOV, K. A., 2012.– *Illiberis (Primilliberis) rotundata* Jordan (Lepidoptera: Zygaenidae: Procridinae) male sex attractant: Optimization and use for seasonal monitoring.– *Entomological Science*, **15**: 137-139.
- SUBCHEV, M., KOSHIO, C., TOSHOVA, T., EFETOV, K. A. & FRANCKE, W., 2013.– (2R)-butyl (7Z)-dodecenoate, a main sex pheromone component of *Illiberis (Primilliberis) pruni* Dyar (Lepidoptera: Zygaenidae: Procridinae)?– *Acta Zoologica Bulgarica*, **65**(3): 391-396.
- SUBCHEV, M., TOSHOVA, T., KOSHIO, C., FRANKE, S., TRÖGER, A., TWELE, R., FRANCKE, W., PICKETT, J. A., WADHAMS, L. J. & WOODCOCK, C. M., 2009.– Identification and biological activity of sex pheromone components from females of the plum moth *Illiberis rotundata* Jordan (Lepidoptera: Zygaenidae: Procridinae).– *Chemoecology*, **19**: 47-54.
- TARMANN, G. M., 1994.– A preliminary review of the classification of the zygaenid subfamily Procridinae (Lepidoptera).– *Nota Lepidopterologica*, Supplement **5**: 115-123.
- TOSHOVA, T. B., SUBCHEV, M. A. & TÓTH, M., 2007.– Role of olfactory and visual stimuli in the mating behaviour of male vine bud moths, *Therapsimima ampellophaga* (Lepidoptera: Zygaenidae).– *European Journal of Entomology*, **104**(1): 57-65.
- WITZGALL, P., KIRSCH, P. & CORK, A., 2010.– Sex pheromones and their impact on pest management.– *Journal of Chemical Ecology*, **36**: 80-100.
- ZAGATTI, P. & RENOU, M., 1984.– Les phéromones sexuelles des zygenes. Le comportement de *Zygaena filipendulae* L. (Lepidoptera, Zygaenidae).– *Annales de la Société entomologique de France*, **20**(4): 439-454.

\*K. A. E.  
Crimean Federal University  
295006 Simferopol  
CRIMEA / *CRIMEA*  
E-mail: efetov.konst@gmail.com

E. E. K.  
Crimean Federal University  
295006 Simferopol  
CRIMEA / *CRIMEA*

E. V. P.  
Crimean Federal University  
295006 Simferopol  
CRIMEA / *CRIMEA*

G. M. T.  
Tiroler Landesmuseum Ferdinandeum  
Naturwissenschaften  
Feldstrasse, 11a  
A-6020 Innsbruck  
AUSTRIA / *AUSTRIA*  
E-mail: g.tarmann@tiroler-landesmuseen.at

\*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 16-V-2015)

(Revisado y aceptado / *Revised and accepted* 25-III-2016)

(Publicado / *Published* 30-IX-2016)